

Optimized Effective Potential for Extended Hubbard Model

I. V. Solovyev

*JRCAT-Angstrom Technology Partnership,
1-1-4 Higashi, Tsukuba, Ibaraki 305-0046, Japan
and Institute of Metal Physics, Russian Academy
of Sciences, Ekaterinburg GSP-170, Russia*

(Dated: February 2, 2008)

Abstract

Antiferromagnetic and charge ordered Hartree-Fock solutions of the one-band Hubbard model with on-site and nearest-neighbor Coulomb repulsions are exactly mapped onto an auxiliary local Kohn-Sham (KS) problem within a density-functional theory. The mapping provides a new insight into the interpretation of the KS equations. (i) With an appropriate choice of the basic variable, there is a universal form of the KS potential, which is applicable both for the antiferromagnetic and the charge ordered solutions. (ii) The Kohn-Sham and Hartree-Fock eigenvalues are interconnected by a scaling transformation. (iii) The band-gap problem is attributed to the derivative discontinuity of the basic variable as the function of the electron number, rather than a finite discontinuity of the KS potential. (iv) It is argued that the conductivity gap and the energies of spin-wave excitations can be entirely defined by the parameters of the ground state and the KS eigenvalues.

PACS numbers: 71.15.Mb, 71.45.Gm, 71.45.Lr, 75.30.Fv

I. INTRODUCTION

Practical success of the Kohn-Sham (KS) scheme of density-functional theory (DFT)^{1,2} in theoretical description of the ground-state properties of molecules and solids lies in the idea of substitution of the real interacting many-electron system with an auxiliary system of noninteracting particles moving in an effective local potential. Similar ideas can be found recently in the dynamic mean-field theory of strongly correlated systems,³ which is based on a mapping of lattice models (e.g., the Hubbard model or the Kondo model) onto a local single-impurity problem. This mapping is well elaborated only for a limited number of examples like the homogeneous electron gas in the density-functional theory [the so-called local-spin-density approximation (LSDA)], or in the limit of infinite lattice coordination in the dynamic mean-field theory. However, many general aspects of such a local noninteracting representation underlying the KS scheme are not well understood. One of the most controversial issues is the physical meaning (if any) of the KS eigenvalues, and their relationship with true physical one-particle excitations.

The major problem, which was intensively discussed in the 1980s,^{4,5,6,7,8,9,10,11,12,13} is the description of fundamental conductivity gaps in semiconductors and insulators. It is generally accepted that the KS one-electron orbital energies have no precise physical meaning, except for the highest occupied eigenvalue, which is equal to the chemical potential (minus the work function). Formally, the conductivity gap is not the ground-state property and includes the excited-state effects. The LSDA systematically underestimate the band gaps. The band-gap problem is typically ascribed to the finite discontinuity of the KS potential as the function of the electron number, meaning that for an extra electron, the KS potential may have an additional constant compared to the KS potential designed for the ground state. In other words, this is equivalent to the statement that the band-gap problem for semiconductors and insulators may be resolved by a rigid upward shift of the conduction bands in the KS picture for the ground state. This procedure is frequently referred to as a "scissor operator".

None of the standard density functionals, such as the LSDA or generalized gradient expansions show such a discontinuity. The origin and the magnitude of this discontinuity is still an open question. Several authors argued that the discontinuity can be large, and the single KS potential (even an exact one) designed for the ground state does not give

the correct gap.^{5,6,7,8} In accordance to their opinion, the solution for the band-gap problem should be sought in the nonlocality and the energy-dependence of the electron self-energy, which is beyond the standard KS formulation, and can be obtained, for example, in the framework of the *GW* approximation.^{7,8,9,13} On the contrary, Gunnarsson and Schönhammer suggested that the actual discontinuity of the exact KS potential may be small, and the main problem lies in the LSDA.¹⁰ Their original arguments were based on a numerical analysis of a Hubbard-type model with the on-site Coulomb repulsion only. However, later they also argued that the intersite Coulomb interaction can produce a large discontinuity of the KS potential.¹¹ The disputed issues have been addressed again recently in Refs. 14 and 15, using two different viewpoints on the formulation of the density functional-theory on a discrete lattice.

In this work we discuss several general aspects of the density-functional theory for the one-band extended Hubbard model. We focus our attention on the analysis of antiferromagnetic (AF) and charge ordered insulating states, the description of which was a traditionally difficult subject for *ab initio* DFT methods.^{16,17,18} From the very beginning we restrict ourselves by the framework of the (static) mean-field (MF) approximation. Physically, this means that we take into the consideration only the (classical) Coulomb and exchange interactions among the electrons, and neglect the correlation effects. In this sense our density functional theory for the Hubbard model is certainly less advanced rather than similar theories existing in the literature and dealing with a formally exact problem numerically.^{10,11,14,15} On the other hand, we hope that such a simplified mean-field analysis may be rather instructive, because we will be able to treat the problem analytically and give a clear picture for several disputed moments. Our primary goal will be to show how the same mean-field problem can be formulated in two different forms, one of which is the conventional Hartree-Fock (HF) theory, where the one-electron orbitals have a clear physical meaning based on Koopmans' theorem, and the other one is the KS scheme in the DFT. The main point, which we would like to emphasize in this work is that the meaning of the KS band structure may not be so formal, as it is currently understood. The relationship between the KS eigenvalues and the true one-particle excitations may be more fundamental, and is not necessarily based only on the discontinuity property of the exact KS potential. We do not discuss here the relative stability of different solutions of the extended Hubbard model. Such an analysis can be found elsewhere (see, e.g., Refs. 19,20,21 and references therein).

The paper is organized as follows: the total energy functional is defined in Sec.II. The KS potential is derived in Sec.III by using the optimized effective potential (OEP) method.²² The connection between the OEP method, the density functional theory and the unrestricted Hartree-Fock approach is discussed in Sections IV and V. In Sec.VI we discuss the connection between the conductivity gap, the energies of spin-wave excitations and the KS one-electron energies. Brief summary will be given in Sec.VII.

II. TOTAL ENERGY

The one-band extended Hubbard Hamiltonian with on-site (U) and nearest-neighbor (V) Coulomb repulsions is defined as

$$\hat{\mathcal{H}} = -t \sum_{\langle ij \rangle} \sum_{\sigma} \left(\hat{c}_i^{\sigma\dagger} \hat{c}_j^{\sigma} + \hat{c}_j^{\sigma\dagger} \hat{c}_i^{\sigma} \right) + U \sum_i \hat{n}_i^{\uparrow} \hat{n}_i^{\downarrow} + V \sum_{\langle ij \rangle} \sum_{\sigma, \sigma'} \hat{n}_i^{\sigma} \hat{n}_j^{\sigma'}, \quad (1)$$

where $\hat{n}_i^{\sigma} = \hat{c}_i^{\sigma\dagger} \hat{c}_i^{\sigma}$ and $\hat{c}_i^{\sigma\dagger}$ ($\hat{c}_i^{\sigma}$) creates (annihilates) an electron on the site i with spin $\sigma = \uparrow$ or $\downarrow$. A nearest-neighbor pair is denoted by $\langle ij \rangle$. The kinetic hopping between nearest-neighbor orbitals is denoted by t . We use the convention, in accordance with which capital letters are reserved for absolute values of parameters of Coulomb interactions and the KS potential, whereas small letters stand for the same parameters in units of $2t$ (for example, $u = U/2t$, etc.).

Let us consider the mean-field solution of the extended Hubbard model, where the ground-state wavefunction of the many-electron system is approximated by a single Slater determinant $|S[\psi_m^{\sigma}]\rangle$ of one-electron orthonormal orbitals $\{\psi_m^{\sigma}\}$. Then, the total energy is given by

$$\mathcal{E}_{\text{MF}}[\psi_m^{\sigma}] = \langle S[\psi_m^{\sigma}] | \hat{\mathcal{H}} | S[\psi_m^{\sigma}] \rangle, \quad (2)$$

and can be expressed in terms of the one-electron density matrix

$$\rho^{\sigma}[i, j] = \sum_m \langle \psi_m^{\sigma} | \hat{c}_i^{\sigma\dagger} \hat{c}_j^{\sigma} | \psi_m^{\sigma} \rangle \quad (3)$$

as $\mathcal{E}_{\text{MF}} = \mathcal{T} + \mathcal{E}_{\text{H}} + \mathcal{E}_{\text{X}}$ (see, e.g., Ref. 2), where $\mathcal{T}$ is the (noninteracting) kinetic energy:

$$\mathcal{T} = -t \sum_{\langle ij \rangle} \sum_{\sigma} (\rho^{\sigma}[i, j] + \rho^{\sigma}[j, i]), \quad (4)$$

$\mathcal{E}_H$ is the Hartree (Coulomb) energy:

$$\mathcal{E}_H = \frac{U}{2} \sum_i (\rho^\uparrow[i, i] + \rho^\downarrow[i, i])^2 + V \sum_{\langle ij \rangle} (\rho^\uparrow[i, i] + \rho^\downarrow[i, i]) (\rho^\uparrow[j, j] + \rho^\downarrow[j, j]), \quad (5)$$

and $\mathcal{E}_X$ is the exchange (Fock) energy:

$$\mathcal{E}_X = -\frac{U}{2} \sum_{i\sigma} (\rho^\sigma[i, i])^2 - V \sum_{\substack{\langle ij \rangle \\ \sigma}} \rho^\sigma[i, j] \rho^\sigma[j, i]. \quad (6)$$

Both $\mathcal{E}_H$ and $\mathcal{E}_X$ include the self-interaction.

Our goal is to simulate a spin or/and charge ordering on the bipartite one-dimensional (1D), two-dimensional (2D), simple cubic (SC) or base-centered cubic (BCC) lattice, for which the spin or/and charge densities are periodically distributed between two types of atoms $\nu = 1, 2$ resulting in the doubling of the unit cell, as it is shown in Fig.1. Thus, the site-indices are specified by the position of the doubled unit cell $\mathbf{R}$ and the type of the site: $i \equiv (\mathbf{R}\nu)$. ρ^σ is the periodic object: $\rho^\sigma[\mathbf{R}'\nu', \mathbf{R}\nu] = \rho^\sigma[(\mathbf{R}' - \mathbf{R})\nu', 0\nu]$, implying that the one-electron orbitals in Eq.(3) are the Bloch states: i.e., $m \equiv (\mathbf{k}\ell)$ and $\hat{c}_{\mathbf{R}\nu}^\sigma |\psi_{\mathbf{k}\ell}^\sigma\rangle = e^{i\mathbf{k}\mathbf{R}} \hat{c}_{0\nu}^\sigma |\psi_{\mathbf{k}\ell}^\sigma\rangle$, where $\mathbf{k}$ is a vector in the first Brillouin zone of the doubled unit cell, and ℓ is the remaining (band) index. In accordance with the definition (1), the range of interactions in the Hamiltonian $\hat{\mathcal{H}}$ is restricted by the nearest neighbors. If we combine this fact with the symmetry properties of the density matrix, i.e., the periodicity and the invariance with respect to the local symmetry operations for the lattices shown in Fig.1, it is rather easy to see that the total energy (2) will be entirely represented by the elements of the density matrix with $\mathbf{R} = \mathbf{R}' = 0$, which can be formally composed into the 2×2 matrix:

$$\hat{\rho}^\sigma = \begin{pmatrix} \rho^\sigma[01, 01] & \rho^\sigma[01, 02] \\ \rho^\sigma[02, 01] & \rho^\sigma[02, 02] \end{pmatrix}. \quad (7)$$

Since the inversion is one of the local symmetry operations, $\hat{\rho}^\sigma$ may be taken in the real symmetric form.

In the following we will focus our attention on the three possible ordered states:

the antiferromagnetic state near to half filling,

the charge-ordered nonmagnetic (CON) state near to half filling,

the charge-ordered ferromagnetic (COF) state near to quarter filling.

The number of electrons in the doubled unit cell is specified as $n_e = n_0 \pm \eta$, where n_0 is the integer electron number at half/quarter filling ($n_0 = 2$ for the AF and CON state, and $n_0 = 1$ for

the COF state) and η describes a small deviation away from half/quarter filling. Then, the elements of the density matrix for these three cases obey the series of symmetry constraints listed in Table I, reducing the number of independent elements in $\hat{\rho}^{\uparrow,\downarrow}$, for a given n_e , to the expectation values of only two pseudospins calculated for the majority-spin channel:

$$\tau_x = \text{Tr}(\hat{\rho}^{\uparrow}\hat{\sigma}_x), \quad \tau_z = \text{Tr}(\hat{\rho}^{\uparrow}\hat{\sigma}_z), \quad (8)$$

where $\hat{\sigma}_x$ and $\hat{\sigma}_z$ are 2×2 Pauli matrices. In the AF state, τ_z is equal to the local magnetic moment. If we define the number of the doubled unit cells as N , and the total energy per doubled unit cell as $E_{\text{MF}} \equiv \mathcal{E}/N = T + E_{\text{H}} + E_{\text{X}}$, we obtain from Eqs.(4)-(6) the following expressions for the kinetic energy (T), the Hartree energy (E_{H}), and the exchange energy (E_{X}) per doubled unit cell, correspondingly

$$T[\tau_x] = -n_0 z t \tau_x, \quad (9)$$

$$E_{\text{H}}[\tau_z] = \frac{1}{4} n_e^2 (U + zV) + \frac{1}{4} (U - zV) (\tau_z + \tau_z^{\downarrow})^2, \quad (10)$$

and

$$E_{\text{X}}[\tau_x, \tau_z] = -\frac{1}{4} n_0 U \left[\left(\frac{n_e}{n_0} \right)^2 + \tau_z^2 \right] - \frac{1}{4} n_0 z V \tau_x^2, \quad (11)$$

where z is the coordination number. In Eq.(10), $\tau_z^{\downarrow}$ is $-\tau_z$, τ_z and 0 for the AF, CON and COF states, respectively. The prefactor n_0 in Eqs.(9) and (11) counts the spin contributions. Due to the symmetry properties of the density matrix (Table I), we have to equivalent contributions to T and E_{X} coming from the majority-spin and minority-spin channels in the case of AF and CON ordering. This is reflected in the prefactor $n_0=2$, which coincides with the number of electrons at half filling. In the case of COF ordering, the minority-spin states are unoccupied. Therefore, we have to count only the majority-spin channel and take $n_0=1$, which again coincides with the number of electrons at quarter filling.

III. OPTIMIZED EFFECTIVE POTENTIAL

There are two possible ways of searching the ground state for the mean-field total energy functional (2). One is the standard Hartree-Fock approach, which will be discussed in Sec.V. Here we start with an alternative provided by the KS scheme and assume that the one-electron orbitals $\{\psi_{\mathbf{k}\ell}^{\sigma}\}$ can be obtained as $n_e \times N$ lowest energy eigenfunctions of the

following Hamiltonians:

$$\hat{\mathcal{H}}_{\text{KS}}^\sigma = -t \sum_{\langle \mathbf{R}1\mathbf{R}'2 \rangle} \left(\hat{c}_{\mathbf{R}1}^{\sigma\dagger} \hat{c}_{\mathbf{R}'2}^\sigma + \hat{c}_{\mathbf{R}'2}^{\sigma\dagger} \hat{c}_{\mathbf{R}1}^\sigma \right) + \sum_{\mathbf{R}} \left(B_1^\sigma \hat{c}_{\mathbf{R}1}^{\sigma\dagger} \hat{c}_{\mathbf{R}1}^\sigma + B_2^\sigma \hat{c}_{\mathbf{R}2}^{\sigma\dagger} \hat{c}_{\mathbf{R}2}^\sigma \right). \quad (12)$$

As in Eq.(1), the notation $\langle \mathbf{R}1\mathbf{R}'2 \rangle$ means that the summation is restricted by the nearest neighbors, which are now of different types (see Fig.1). If $\mathbf{r}$ is the relative position of two different atoms within one unit cell, the notation $\langle \mathbf{R}1\mathbf{R}'2 \rangle$ is equivalent to the condition $|\mathbf{R}' + \mathbf{r} - \mathbf{R}| = |\mathbf{r}|$. Thus, $\hat{\mathcal{H}}_{\text{KS}}^{\uparrow,\downarrow}$ retains the kinetic part of by the original Hamiltonian $\hat{\mathcal{H}}$. The interacting part is substituted by the effective one-electron KS potentials $\{B_\nu^\sigma\}$, which are assumed to be site-diagonal ("local" with respect to the site-indices), do not depend on $\mathbf{R}$ (as it is requested by the periodicity of the antiferromagnetic and charge-ordered states), but may depend on the type of the atom ν . $\{B_\nu^\sigma\}$ can be found in the spirit of optimized effective potential method^{22,26} as the solution of the variational problem (for the given electron number n_e):

$$\frac{\partial}{\partial B_\nu^\sigma} \left(T[\tau_x] + E_{\text{H}}[\tau_z] + E_{\text{X}}[\tau_x, \tau_z] - \sum_{\lambda} \lambda \mathcal{F}_\lambda[\{B_\nu^\sigma\}] \right) \Big|_{n_e} = 0, \quad (13)$$

provided that $\{B_\nu^\sigma\}$ obey the series of constraints $\mathcal{F}_\lambda[\{B_\nu^\sigma\}]$, which guarantee the required symmetry of the density matrix (Table I). Thus, $\{B_\nu^\sigma\}$ can be interpreted as the exact KS potentials for exchange alone.² The Hamiltonian (12) can be diagonalized in the basis of Bloch orbitals

$$\begin{pmatrix} \hat{c}_1^{\sigma\dagger}(\mathbf{k}) \\ \hat{c}_2^{\sigma\dagger}(\mathbf{k}) \end{pmatrix} = \frac{1}{\sqrt{N}} \sum_{\mathbf{R}} e^{i\mathbf{k}\mathbf{R}} \begin{pmatrix} \hat{c}_{\mathbf{R}1}^{\sigma\dagger} \\ e^{i\mathbf{k}\mathbf{r}} \hat{c}_{\mathbf{R}2}^{\sigma\dagger} \end{pmatrix}. \quad (14)$$

The eigenfunctions of $\hat{\mathcal{H}}_{\text{KS}}^\sigma$ are

$$\begin{pmatrix} |\psi_{\mathbf{k}-}^\sigma\rangle \\ |\psi_{\mathbf{k}+}^\sigma\rangle \end{pmatrix} = \begin{pmatrix} \cos \vartheta_{\mathbf{k}}^\sigma & \sin \vartheta_{\mathbf{k}}^\sigma \\ -\sin \vartheta_{\mathbf{k}}^\sigma & \cos \vartheta_{\mathbf{k}}^\sigma \end{pmatrix} \begin{pmatrix} \hat{c}_1^{\sigma\dagger}(\mathbf{k})| \rangle \\ \hat{c}_2^{\sigma\dagger}(\mathbf{k})| \rangle \end{pmatrix}, \quad (15)$$

where $| \rangle$ denotes the vacuum state with no electrons in it.

We note that due to the symmetry constraints (Table I), we have to consider only the case $-B_1^\uparrow = B_2^\uparrow \equiv B$. Then, the eigenvalues of $\hat{H}_{\text{KS}}^\uparrow$ are

$$\varepsilon_{\pm}^\uparrow(\mathbf{k}) = \pm \sqrt{B^2 + 4t^2 f^2(\mathbf{k})} \quad (16)$$

and the angle $\vartheta_{\mathbf{k}}^\uparrow$ in Eq.(15) is given by

$$\text{tg} \vartheta_{\mathbf{k}}^\uparrow = \frac{f(\mathbf{k})}{b + \sqrt{b^2 + f^2(\mathbf{k})}}, \quad (17)$$

where

$$f(\mathbf{k}) = \frac{1}{2} \sum_{\mathbf{R}: |\mathbf{r}-\mathbf{R}|=|\mathbf{r}|} \cos \mathbf{k}(\mathbf{r}-\mathbf{R})$$

is the structure factor. [The explicit form of $f(\mathbf{k})$ for different lattices is shown in Fig.1]. Then, putting the wavefunctions (15) in equations (3), (7) and (8), we obtain after some algebra:

$$\tau_x = \frac{2}{zN} \sum_{\mathbf{k}\ell} \frac{(-\ell)f^2(\mathbf{k})}{\sqrt{b^2 + f^2(\mathbf{k})}} \Theta[\varepsilon_F - \varepsilon_\ell^\uparrow(\mathbf{k})], \quad (18)$$

$$\tau_z = \frac{1}{N} \sum_{\mathbf{k}\ell} \frac{(-\ell)b}{\sqrt{b^2 + f^2(\mathbf{k})}} \Theta[\varepsilon_F - \varepsilon_\ell^\uparrow(\mathbf{k})], \quad (19)$$

where $\ell=\pm 1$, and the Fermi energy ε_F is determined by the condition:

$$n_0 \sum_{\mathbf{k}\ell} \Theta[\varepsilon_F - \varepsilon_\ell^\uparrow(\mathbf{k})] = n_e N. \quad (20)$$

Then, it is rather straightforward to show for any fixed electron number n_e that

$$\left. \frac{\partial \tau_x}{\partial b} \right|_{n_e} = -\frac{2b}{z} \left. \frac{\partial \tau_z}{\partial b} \right|_{n_e}. \quad (21)$$

Therefore, τ_x and τ_z are not independent variables and the variational procedure of the OEP approach is equivalent to a DFT in which τ_z participates as the basic variable. Indeed, using Eqs.(9) and (21) we obtain $(\partial T[\tau_x]/\partial \tau_z)|_{n_e=n_0} B$. As it was discussed earlier, the prefactor n_0 counts the spin channels involved into the problem. Then, variational principle (13) leads to the standard definition for the KS potential conjugated with the basic variable τ_z :

$$B = -\frac{1}{n_0} \left\{ \left. \frac{\partial E_H[\tau_z]}{\partial \tau_z} \right|_{n_e} + \left. \frac{\partial E_X[\tau_x, \tau_z]}{\partial \tau_z} \right|_{n_e} \right\}, \quad (22)$$

where the minus sign is related with our definition of τ_z and B : for $\sigma=\uparrow$, the solution with $\tau_z=\rho^\uparrow[01,01]-\rho^\uparrow[02,02]>0$ corresponds to the potential $B_1^\uparrow=-B_2^\uparrow\equiv-B<0$. The exchange part in Eq.(22) is given by

$$B_X \equiv -\frac{1}{n_0} \left. \frac{\partial E_X[\tau_x, \tau_z]}{\partial \tau_z} \right|_{n_e} = \frac{1}{2} U \tau_z - V \tau_x b \quad (23)$$

[see Eqs.(11) and (21)]. The Coulomb part $B_H \equiv (-1/n_0)(\partial E_H[\tau_z]/\partial \tau_z)|_{n_e}$ is 0, $(U-zV)\tau_z$ and $\frac{1}{2}(U-zV)\tau_z$ for the AF, CON and COF states, respectively [see Eq.(10)]. Then, Eq.(22) can be transformed to the self-consistent equation for the KS potential $b=b_H+b_X$:

$$b = \frac{1}{2} u_{\text{HF}} \tau_z - v \tau_x b, \quad (24)$$

where u_{HF} is u , $2zv - u$ and zv for the AF, CON and COF states, respectively. Eq.(24) further yields:

$$b = \frac{1}{2}u_{\text{KS}}[\tau_{\text{x}}]\tau_{\text{z}}, \quad (25)$$

where the effective coupling $u_{\text{KS}}[\tau_{\text{x}}]$ is given by

$$u_{\text{KS}}[\tau_{\text{x}}] = \frac{u_{\text{HF}}}{1 + v\tau_{\text{x}}}. \quad (26)$$

Putting Eq.(25) in Eq.(23), one can find for the self-consistent exchange potential b_{x} :

$$b_{\text{x}} = \frac{1}{2} \left[\frac{v\tau_{\text{x}}}{1 + v\tau_{\text{x}}} u_{\text{HF}} - u \right] \tau_{\text{z}}. \quad (27)$$

The local KS potential has the same form [Eq.(25)] for different ordered states, although the driving force for the ordering may be different: u in the AF state and v in the CO states. The local form of the KS potential for the CO states has one important implication which we would like to emphasize here. It is well established that Hubbard-type models with solely on-site Coulomb interactions treated on the level of mean-field theories may exhibit the charge-ordered solutions (see, e.g., Ref.23). The present work suggests that such "on-site Coulomb interactions" may be interpreted in the Kohn-Sham sense, as parameters of an auxiliary local KS problem, whose real physical meaning may be different and implies the contributions of longer-range Coulomb interactions.

u_{KS} has the common renormalization factor $(1+v\tau_{\text{x}})^{-1}$ which is driven by the intersite Coulomb interaction v and depends on τ_{z} through τ_{x} . The dependence $\tau_{\text{x}}[\tau_{\text{z}}]$ is shown in Fig.2 for several lattices and $n_e=n_0$. We note: (i) for a given τ_{z} , τ_{x} is larger when z is smaller. Thus, the renormalization is the most important for 1D systems. (ii) In the framework of our model, $\tau_{\text{z}}=0$ may be regarded as a homogeneous solution.¹⁶ Since τ_{x} is the decreasing function of τ_{z} , we have $u_{\text{KS}}[0] \leq u_{\text{KS}}[\tau_{\text{z}}]$ meaning that the parameter u_{KS} being once evaluated in the homogeneous limit and employed afterwards for the whole range of τ_{z} is generally "overscreened". It is therefore clear that an attempt to substitute the real effective coupling $u_{\text{KS}}[\tau_{\text{z}}]$ with its homogeneous counterpart $u_{\text{KS}}[0]$ in the KS potential will generally underestimate the tendencies towards the antiferromagnetism and the charge ordering, as it is well established for the LSDA.¹⁷ The analytical estimate for $\tau_{\text{z}}=0$ and $n_e=n_0$ is $\tau_{\text{x}}[0]=(2/\pi)^{\log_2 z}$ (except the SC lattice). At the opposite end $\tau_{\text{z}}=1$, corresponding to the limit $u_{\text{KS}}[1] \rightarrow \infty$, we have $\tau_{\text{x}} \rightarrow 1/2b \rightarrow 1/u_{\text{KS}}[1]$, see Eqs.(18) and (25). Then, from

Eq.(26) we obtain $u_{\text{KS}}[1]=u_{\text{HF}}(1+v/u_{\text{KS}}[1])^{-1}$, whence it follows that

$$u_{\text{KS}}[1] = u_{\text{HF}} - v. \quad (28)$$

IV. THE LEVY CONSTRAINED-SEARCH FORMULATION

A very elegant formulation of the density-functional theory was introduced by Levy.² For our purposes it is read as follows: if τ_z is the basic variable, the total energy minimization (for a given electron number) can be performed in two steps:

$$\mathcal{E} = \min_{\tau_z} \left\{ \min_{|\Psi\rangle \rightarrow \tau_z} \langle \Psi | \hat{\mathcal{H}} | \Psi \rangle \right\}. \quad (29)$$

The inner minimization is constrained to all many-electron wavefunctions $|\Psi\rangle$ that give τ_z , while the outer minimization releases the constraint by searching all τ_z . In the mean-filed approximation, the search of all possible many-electron wavefunctions is restricted within the subspace of Slater determinants $|S[\psi_m^\sigma]\rangle$. Thus, $|\Psi\rangle$ is additionally constrained, and we have $\mathcal{E} \leq \mathcal{E}_{\text{MF}}$. The purpose of this section is to show that in all other respects, the OEP approach employed in the previous section is equivalent to the Levy constrained-search formulation (29). The proof is simple and can be carried out along the general line taken in the standard DFT formulation.² Taking into account the explicit form of $\langle S[\psi_m^\sigma] | \hat{\mathcal{H}} | S[\psi_m^\sigma] \rangle$, Eqs.(9)-(11), the inner step in Eq.(29) is equivalent to the search of the one-electron orbitals $\{\psi_m^\sigma\}$ which maximize the expectation value of the pseudospin τ_x for the given τ_z , or equivalently, minimize the (noninteracting) kinetic energy T for the given τ_z . Using the definitions of T and τ_z in terms of the one-particle orbitals [see Eqs. (3), (4), (7) and (8)], this gives the functional

$$\Omega[\psi_m^\uparrow] = -t \sum_m \sum_{\langle \mathbf{R}1\mathbf{R}'2 \rangle} \langle \psi_m^\uparrow | \hat{c}_{\mathbf{R}1}^\dagger \hat{c}_{\mathbf{R}'2}^\dagger + \hat{c}_{\mathbf{R}'2}^\dagger \hat{c}_{\mathbf{R}1}^\dagger | \psi_m^\uparrow \rangle - \lambda^\uparrow \sum_m \sum_{\mathbf{R}} \left(\langle \psi_m^\uparrow | \hat{c}_{\mathbf{R}1}^\dagger \hat{c}_{\mathbf{R}1}^\dagger - \hat{c}_{\mathbf{R}2}^\dagger \hat{c}_{\mathbf{R}2}^\dagger | \psi_m^\uparrow \rangle - \tau_z \right) - \sum_{mm'} \varepsilon_{mm'}^\uparrow \left(\langle \psi_m^\uparrow | \psi_{m'}^\uparrow \rangle - \delta_{mm'} \right) \quad (30)$$

and the minimization condition $\delta\Omega[\psi_m^\uparrow]/\delta\langle\psi_m^\uparrow| = 0$. (The Lagrange multipliers $\varepsilon_{mm'}^\uparrow$ stand for the orthonormalization condition between $|\psi_m^\uparrow\rangle$ and $|\psi_{m'}^\uparrow\rangle$). These result in the equation

$$\hat{h}^\uparrow |\psi_m^\uparrow\rangle = \sum_{m'} \varepsilon_{mm'}^\uparrow |\psi_{m'}^\uparrow\rangle, \quad (31)$$

where

$$\hat{h}^\uparrow = -t \sum_{\langle \mathbf{R}1\mathbf{R}'2 \rangle} \left(\hat{c}_{\mathbf{R}1}^\dagger \hat{c}_{\mathbf{R}'2}^\dagger + \hat{c}_{\mathbf{R}'2}^\dagger \hat{c}_{\mathbf{R}1}^\dagger \right) - \lambda^\uparrow \sum_{\mathbf{R}} \left(\hat{c}_{\mathbf{R}1}^\dagger \hat{c}_{\mathbf{R}1}^\dagger - \hat{c}_{\mathbf{R}2}^\dagger \hat{c}_{\mathbf{R}2}^\dagger \right). \quad (32)$$

Similar equations can be obtained for $\sigma=\downarrow$ by using the symmetry properties of the density matrix from Table I. The Hamiltonian (32) has the same form as the KS Hamiltonian (12). (Note also that since $\hat{h}^\uparrow$ is Hermitian, $\varepsilon_{mm'}^\uparrow$ can be taken in the diagonal form). If τ_z is the expectation value of the pseudospin corresponding to the ground state, the Lagrange multiplier $\lambda^\uparrow$ is equal to the self-consistent potential B . Thus, the inner minimization of the Levy constrained-search formulation is equivalent to the diagonalization of the KS Hamiltonian in the OEP approach.

The conclusion is very natural and, in principle, directly follows from the definition of the KS scheme.^{1,2} Indeed, the mean-field energy $\mathcal{E}_{\text{MF}}$ is determined by only two parameters, τ_x and τ_z . Both of them are requested to come out exactly of the one-electron KS orbitals $\{\psi_m^\sigma\}$: τ_z as the basic variable, and τ_x as the only variable which determines the noninteracting kinetic energy.

V. UNRESTRICTED HARTREE-FOCK APPROACH

In order to clarify the origin of the renormalization factor $(1+v\tau_x)^{-1}$ in the KS potential (25), let us now turn to the Hartree-Fock solution of the extended Hubbard model, where the basic variables are the one-electron orbitals $\{\psi_m^\sigma\}$: $\hat{\mathcal{H}}_{\text{HF}}^\sigma|\psi_m^\sigma\rangle=\delta\mathcal{E}_{\text{MF}}/\delta\langle\psi_m^\sigma|$. Taking into account the periodicity of the ordered states, this gives the Hamiltonian (the uniform potential shift is dropped):

$$\hat{\mathcal{H}}_{\text{HF}}^\uparrow = -t(1+v\tau_x) \sum_{\langle\mathbf{R}|\mathbf{R}'/2\rangle} \left(\hat{c}_{\mathbf{R}1}^\dagger \hat{c}_{\mathbf{R}'2}^\dagger + \hat{c}_{\mathbf{R}'2}^\dagger \hat{c}_{\mathbf{R}1}^\dagger \right) - \frac{1}{2} U_{\text{HF}} \tau_z \sum_{\mathbf{R}} \left(\hat{c}_{\mathbf{R}1}^\dagger \hat{c}_{\mathbf{R}1}^\dagger - \hat{c}_{\mathbf{R}2}^\dagger \hat{c}_{\mathbf{R}2}^\dagger \right) \quad (33)$$

with the parameters of site-diagonal interactions U_{HF} defined earlier. The prefactor $(1+v\tau_x)$ in the off-diagonal part of the HF Hamiltonian comes from the nonlocality of the Fock exchange potential, which is combined with the kinetic term. Since the KS potential is requested to be local, the same effect enters the KS formulation through the renormalization of the site-diagonal interactions, as it is indeed manifested in Eq.(26). Therefore, the terminology "the local potential" and "the local approximation" used in the density-functional theory does not necessarily mean that the nonlocal exchange interactions are totally neglected. They may contribute to the renormalization of the local interactions.

In the framework of the one-band extended Hubbard model, we obtain the trivial connection between the KS and HF Hamiltonians: $\hat{\mathcal{H}}_{\text{KS}}^\uparrow=(1+v\tau_x)^{-1}\hat{\mathcal{H}}_{\text{HF}}^\uparrow$, meaning that they have

the same set of eigenfunctions, but different eigenvalues. The KS band structure is simply the *scaled* HF one. Since $\tau_x \geq 0$ [see Eq.(18)], the scaling is in fact a *contraction*, unless $v=0$. Although such a trivial scaling scenario is a crude simplification of the situation realized in realistic calculations, it does explain the main tendency: not only the band gaps, but also the bandwidths appear to be smaller in the KS picture.²⁴ This is definitely different from the rigid upward shift of the conduction bands, underlying the "scissor operator", which affects only the gap width.

VI. ONE-PARTICLE EXCITATIONS

In this section we establish several useful connections between Kohn-Sham eigenvalues and true one-particle excitation energies for the extended Hubbard Hamiltonian in the mean-field approach. First, we will consider the conductivity gap as an example of excitations for which the total number of particles is not conserved. We will show that although in this case the physical gap can be obtained from the total energy functional and is entirely defined by the parameters of the ground state, the one-electron KS band structure plays a formally fictitious role, in the sense that the KS gap is generally different from the true conductivity gap. Second, we will turn to the analysis of virtual spin-wave excitations in the antiferromagnetic insulating state at half filling and argue that these excitations can be interpreted *directly* in terms of the KS eigenvalues.

A. Conductivity gap

If $n_e = n_0$, the KS scheme yields the insulating solution. The band gap $\Delta_{\text{KS}} = U_{\text{KS}} \tau_z$ is direct and opens in the points $\{\mathbf{k}_0\}$ for which $f(\mathbf{k}_0) = 0$ and $\vartheta_{\mathbf{k}_0}^\dagger = 0$ [see Eqs. (16) and (17)]. In order to calculate the physical conductivity gap, we first take the standard procedure²⁵ and employ the fact that two Hamiltonians, $\hat{\mathcal{H}}_{\text{KS}}^\dagger$ and $\hat{\mathcal{H}}_{\text{HF}}^\dagger$, have the same set of eigenfunctions. Then, the conductivity gap is given by

$$\Delta = \Delta_{\text{KS}} + \langle \psi_{\mathbf{k}_0+}^\dagger | \hat{\mathcal{H}}_{\text{HF}}^\dagger - \hat{\mathcal{H}}_{\text{KS}}^\dagger | \psi_{\mathbf{k}_0+}^\dagger \rangle - \langle \psi_{\mathbf{k}_0-}^\dagger | \hat{\mathcal{H}}_{\text{HF}}^\dagger - \hat{\mathcal{H}}_{\text{KS}}^\dagger | \psi_{\mathbf{k}_0-}^\dagger \rangle. \quad (34)$$

Taking this as an exercise and using Eq.(15) for $\vartheta_{\mathbf{k}_0}^\dagger = 0$, it is very easy to verify that $\Delta = U_{\text{HF}} \tau_z \equiv \Delta_{\text{HF}}$, which is nothing but the consequence of well-known Koopmans' theorem

for the eigenvalues of the HF Hamiltonian (33). (Note, that the correction to Koopmans' theorem due to the relaxation of HF levels vanishes for extensive systems when $N \rightarrow \infty$).

The central question is how the physical gap is related with the ground-state properties, and whether it can be entirely described by the parameters of the ground state and the KS eigenvalues. As it was discussed in the introduction, the most prominent scenario is based on the idea of discontinuity of the KS potential. In the case of finite species the similar problem was investigated for the chemical hardness (half of the band gap in the limit $N \rightarrow \infty$). It was argued that the exact KS potential exhibits a constant jump (discontinuity) when the number of electrons changes through an integer value, and the chemical hardness can be found as the difference of highest occupied KS eigenvalues calculated for the initial and the final electron configurations (see, e.g., Refs. 4,26,27). The constant jump of the potential, however, is not related with the ground-state properties. Similar arguments can be applied for infinite systems, where the band gap can be expressed through the change of the chemical potential.⁸ In the context of the present work, this would be equivalent to the claim that the uniform part of the KS potential (C), which was irrelevant to the search of the ground state and dropped out the analysis in Sections III-V, should exhibit the finite jump across the band gap: $C|_{n_0+\eta} - C|_{n_0-\eta} = \Delta_{\text{HF}} - \Delta_{\text{KS}}$, and the band-gap problem might be resolved via the scissor-type correction.

Below we present a different view on the band-gap problem. To begin with we note that the quantities τ_x , τ_z and b , which do determine the ground state, exhibit no finite discontinuity as the function of the electron number (Fig.3). The analysis presented in preceding sections clearly shows that the physical gap Δ_{HF} can be expressed in terms of these characteristics of the ground state by using another, scaling transformation $\Delta_{\text{HF}} = (1 + v\tau_x)\Delta_{\text{KS}}$.²⁸ Thus, *there is* a unique connection between Δ_{HF} and the ground-state properties. In order to gain more insight into the problem, we argue below that the physical gap Δ_{HF} can be naturally derived from the mean-field total energy functional $\mathcal{E}_{\text{MF}}[\tau_x, \tau_z]$, not exploiting the idea of the finite discontinuity of the exact KS potential. Instead, we argue that the band-gap problem is related with the derivative discontinuity of the basic variable τ_z across the band gap. We start with the general definition:

$$\Delta = \mathcal{E}_{\text{MF}}(n_0N + 1) + \mathcal{E}_{\text{MF}}(n_0N - 1) - 2\mathcal{E}_{\text{MF}}(n_0N). \quad (35)$$

As in Eq.(34), we assume that the addition or the removal of an electron does not destroy

the symmetry of the ground state. (For example, if the ground state of the $n_0 N$ electron system is antiferromagnetic, we assume that the system of $n_0 N \pm 1$ electrons will remain in the antiferromagnetic state, though this requirement may formally contradict to the Nagaoka theorem²⁹). Then, in the limit $N \rightarrow \infty$, Eq.(35) becomes

$$\Delta = \lim_{\eta \rightarrow 0} \left\{ \frac{dE_{\text{MF}}[\tau_x, \tau_z]}{dn_e} \Big|_{n_0 + \eta} - \frac{dE_{\text{MF}}[\tau_x, \tau_z]}{dn_e} \Big|_{n_0 - \eta} \right\}, \quad (36)$$

where $\eta = 1/N$. Taking into account the explicit form of τ_x and τ_z , Eqs. (18) and (19), we will have two contributions to $(dE_{\text{MF}}[\tau_x, \tau_z]/dn_e)|_{n_0 \pm \eta}$: one is due to the change of the band filling with the fixed potential b (the derivative of the Θ -function), and the other one is due to the n_e -dependence of the self-consistent potential itself. Therefore, we can write:

$$\frac{dE_{\text{MF}}[\tau_x, \tau_z]}{dn_e} \Big|_{n_0 \pm \eta} = \frac{\partial E_{\text{MF}}[\tau_x, \tau_z]}{\partial n_e} \Big|_{b, n_0 \pm \eta} + \frac{\partial E_{\text{MF}}[\tau_x, \tau_z]}{\partial b} \Big|_{n_0} \frac{\partial b}{\partial n_e} \Big|_{n_0 \pm \eta}.$$

However, in the ground state it holds $(\partial E_{\text{MF}}[\tau_x, \tau_z]/\partial b)|_{n_0} = 0$ [see Eq.(13)], and we find:

$$\Delta = \lim_{\eta \rightarrow 0} \left\{ \frac{\partial E_{\text{MF}}[\tau_x, \tau_z]}{\partial n_e} \Big|_{b, n_0 + \eta} - \frac{\partial E_{\text{MF}}[\tau_x, \tau_z]}{\partial n_e} \Big|_{b, n_0 - \eta} \right\}. \quad (37)$$

Then, we have the following identity: $(\partial \Theta[\varepsilon_F - \varepsilon_{\pm}^{\uparrow}(\mathbf{k})]/\partial n_e)|_{b, n_0 \pm \eta} = \delta[\varepsilon_F|_{b, n_0 \pm \eta} - \varepsilon_{\pm}^{\uparrow}(\mathbf{k})](\partial \varepsilon_F/\partial n_e)|_{b, n_0 \pm \eta}$.

In the limit $\eta \rightarrow 0$, the δ -function will pick up only those points in the Brillouin zone which correspond to the band-gap edges, i.e., $\mathbf{k} \in \{\mathbf{k}_0\}$. Therefore, we obtain from Eq.(20):

$$\lim_{\eta \rightarrow 0} \sum_{\mathbf{k}} \frac{\partial \Theta[\varepsilon_F - \varepsilon_{\pm}^{\uparrow}(\mathbf{k})]}{\partial n_e} \Big|_{b, n_0 \pm \eta} \equiv \lim_{\eta \rightarrow 0} \sum_{\mathbf{k} \in \{\mathbf{k}_0\}} \frac{\partial \Theta[\varepsilon_F - \varepsilon_{\pm}^{\uparrow}(\mathbf{k})]}{\partial n_e} \Big|_{b, n_0 \pm \eta} = \frac{N}{n_0}.$$

This immediately yields: $\lim_{\eta \rightarrow 0} (\partial \tau_x/\partial n_e)|_{b, n_0 \pm \eta} = 0$ [see Eq.(18) and note that $f(\mathbf{k}_0) = 0$], meaning that there is no derivative discontinuity of the kinetic energy (9) across the band gap caused by the change of the band filling with the fixed KS potential. (Do not mix it with the behavior of τ_x in Fig.3, which shows the total effect, including the change of the KS potential b).

On the contrary, the derivative of the basic variable τ_z shows the discontinuity: $n_0 \lim_{\eta \rightarrow 0} (\partial \tau_z/\partial n_e)|_{b, n_0 \pm \eta} = \mp 1$. Then, putting these results in Eq.(37) and using the explicit form of $E_{\text{MF}}[\tau_x, \tau_z]$, Eqs.(9)-(11), we again arrive at Koopmans' theorem for the HF gap: $\Delta = U_{\text{HF}} \tau_z \equiv \Delta_{\text{HF}}$.

Thus, although the KS gap is generally different from the true conductivity gap, Δ_{HF} in our model, the latter can still be derived from the total energy functional $E_{\text{MF}}[\tau_x, \tau_z]$ using the definition (37).

B. Virtual spin-wave excitations

Let us consider the energies of spin-wave excitations in the antiferromagnetic insulating state at half filling.

For these purposes it is convenient to work with the local representation for the one-electron potential provided by the KS scheme. Then, the total energy changes caused by small nonuniform rotations of the spin magnetic moments near the (antiferromagnetic) equilibrium, i.e., the processes relevant to the spin-wave excitations, can be exactly mapped onto the Heisenberg-type model^{30,31,32}

$$E[\{\mathbf{e}_i\}] = -\frac{1}{2} \sum_{ij} J_{ij} \mathbf{e}_i \cdot \mathbf{e}_j, \quad (38)$$

where $\mathbf{e}_i$ is the direction of the spin magnetic moment at the site i . The parameters of the model can be expressed in terms of the one-electron Green function $G_{ij}^{\uparrow,\downarrow}(\varepsilon)$ constructed for the KS Hamiltonian (12) as^{30,31}

$$J_{ij} = \frac{\Delta_{\text{ex}}^i \Delta_{\text{ex}}^j}{2\pi} \text{Im} \int_{-\infty}^{\varepsilon_F} d\varepsilon G_{ij}^{\uparrow}(\varepsilon) G_{ji}^{\downarrow}(\varepsilon), \quad (39)$$

where $\Delta_{\text{ex}}^i \equiv B_{\nu}^{\downarrow} - B_{\nu}^{\uparrow}$ is the local exchange splitting at the site $i \equiv (\mathbf{R}\nu)$. The mapping is based on so-called local force theorem. One can argue rather generally that if the spin-magnetization density ($\pm\tau_z$ in our model for the antiferromagnetic state) is the basic variable in a (spin)-density-functional theory, and is requested to come out exactly of the one-electron KS orbitals, such a mapping onto the Heisenberg-type model should be exact.³²

For two nonequivalent sites of our model we have: $\Delta_{\text{ex}}^{\mathbf{R}1} = -\Delta_{\text{ex}}^{\mathbf{R}2} = U_{\text{KS}}\tau_z$. Thus, the strength of the interatomic exchange couplings J_{ij} is directly related with the magnitude of the KS gap $\Delta_{\text{KS}} = U_{\text{KS}}\tau_z$.

Let us consider the large- u_{KS} limit. In the lowest order of the $1/u_{\text{KS}}$ expansion, the interatomic magnetic interactions are restricted by the nearest neighbors $\langle ij \rangle$. Corresponding elements of the Green function are given by $G_{\langle ij \rangle}^{\uparrow}(\varepsilon) = G_{\langle ji \rangle}^{\downarrow}(\varepsilon) \simeq -t/(\varepsilon^2 - \frac{1}{4}U_{\text{KS}}^2\tau_z^2)$. Putting them in Eq.(39) and noting that in the large- u_{KS} limit $\tau_z=1$ and $U_{\text{KS}}=U-V$ [see Eq.(28)], we arrive at the well-known result for the effective magnetic coupling between nearest neighbors in the extended Hubbard model:^{19,20,21,33} $J_{\langle ij \rangle} = t^2/(U-V)$, or $J_{\langle ij \rangle} = t^2/\Delta_{\text{KS}}$. [Note, that the sign of $J_{\langle ij \rangle}$ depends on the convention used for the directions of the spin magnetic moments in Eq.(38). If we refer to the antiferromagnetic ground state, $\mathbf{e}_i \cdot \mathbf{e}_j = -1$, and therefore

$J_{\langle ij \rangle} > 0$]. Thus, the interatomic magnetic interactions can be regarded as virtual hoppings between effective KS levels, and in this sense the KS band structure has the clear physical meaning.

Finally, we would like to mention one rather interesting implication of two examples discussed in this section to the theory of strongly-correlated materials. The fact that the intersite Coulomb interaction V "renormalizes" primarily the exchange coupling $J_{\langle ij \rangle}$ in the antiferromagnetic state, but not the conductivity gap was well-known from earlier studies of the extended Hubbard model. Such an apparently different influence of the intersite Coulomb interaction V on different physical quantities has led to some ambiguity in the determination of the physical parameter, which controls the strength of correlation effects in solids (see, e.g., discussions in Refs.20,21). The present work suggests that these two properties are not necessarily independent. In the mean-field approach, such a behavior means that there are (at least) two different representations for the one-particle spectrum, which correspond to the same ground state. One representation is essentially nonlocal, but allows the spectroscopic interpretation based on Koopmans' theorem (the Hartree-Fock scheme). Another representation is local and can be directly applied to the analysis of interatomic magnetic interactions (the Kohn-Sham scheme).

VII. SUMMARY

Being based on the one-band extended Hubbard model, we have discussed several aspects of the Kohn-Sham formulation of the density-functional theory in solids. Although the conclusions are strictly applicable only to the mean-field solutions of this particular model, the analysis presented in the paper suggests that the old-standing dilemma between the description of the ground-state properties in the density-functional theory and the description of (at least some) one-particle excitations may be resolved in a more rational way. Particularly, we have shown that the conductivity gap and the interatomic magnetic interactions relevant to the spin-wave excitations for this model are entirely defined by the parameters of the ground state. This therefore means that in this case there is no fundamental difference between the ground-state properties, the conductivity gap and the spin-wave excitation energies, and all of them can be described by the same set of one-electron Kohn-Sham equations. The

KS eigenvalues may not be necessarily coincide with the true one-particle excitation energies, depending on the form in which the nonlocal exchange interactions contribute to each particular type of excitations.

Thus, we hope that results of our model analysis may provide a new insight into the interpretation of one-electron equations underlying the Kohn-Sham scheme and stimulate new researches in this direction. As an extension of the present analysis, we believe that it would be particularly interesting: (i) to go beyond the mean-field approximation and to take into account the correlation effects; (ii) to go beyond the one-band picture, to take into account the effects of orbital degeneracy and to clarify the roles played by intersite Coulomb interactions in the problem of orbital polarization in solids.³⁴

Acknowledgments

I thank K. Terakura and F. Aryasetiawan for useful discussions. The present work is partly supported by New Energy and Industrial Technology Development Organization (NEDO).

-
- ¹ W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
 - ² For review see, e.g., R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules* (Oxford Univ. Press, New York, 1989).
 - ³ A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).
 - ⁴ J. P. Perdew, R. G. Parr, M. Levy, and J. L. Balduz, Phys. Rev. Lett. **49**, 1691 (1982).
 - ⁵ J. P. Perdew and M. Levy, Phys. Rev. Lett. **51**, 1884 (1983).
 - ⁶ L. J. Sham and M. Schlüter, Phys. Rev. Lett. **51**, 1888 (1983).
 - ⁷ R. W. Godby, M. Schlüter, and L. J. Sham, Phys. Rev. Lett. **56**, 2415 (1986).
 - ⁸ R. W. Godby, M. Schlüter, and L. J. Sham, Phys. Rev. B **37**, 10159 (1988).
 - ⁹ M. S. Hybertsen and S. G. Louie, Phys. Rev. B **34**, 5390 (1986).
 - ¹⁰ O. Gunnarsson and K. Schönhammer, Phys. Rev. Lett. **56**, 1968 (1986).
 - ¹¹ K. Schönhammer and O. Gunnarsson, J. Phys. C: Solid State Phys. **20**, 3675 (1987).

- ¹² L. J. Sham and M. Schlüter, Phys. Rev. Lett. **60**, 1582 (1988); O. Gunnarsson and K. Schönhammer, *ibid.* **60**, 1583 (1988).
- ¹³ For review see, e.g., R. O. Jones and O. Gunnarsson, Rev. Mod. Phys. **61**, 689 (1989); F. Aryasetiawan and O. Gunnarsson, Rep. Prog. Phys. **61**, 237 (1998).
- ¹⁴ A. Schindlmayr and R. W. Godby, Phys. Rev. B **51**, 10427 (1995).
- ¹⁵ K. Schönhammer, O. Gunnarsson, and R. M. Noack, Phys. Rev. B **52**, 2504 (1995).
- ¹⁶ Note, that the antiferromagnetism or the charge ordering implies the existence of magnetic or charge inhomogeneities on the interatomic scale. Most part of the first-principles DFT methods is based either on the purely homogeneous electron gas picture for the exchange-correlation energy (LSDA), or takes into account only local effects of the inhomogeneity through gradient expansions^{1,2}.
- ¹⁷ A canonical example of the systems where LSDA fails to reproduce the correct AF ground state is the two-dimensional cuprates [W. E. Pickett, Rev. Mod. Phys. **61**, 433 (1989)]. The charge-ordered systems are less studied: the problem is complicated by the fact that the charge ordering often exists only in a combination with a peculiar spin and orbital ordering, leading to the necessity of simultaneous correct account for *all* three degrees of freedom. An example of such a failure of the LSDA is Fe₃O₄ [Z. Zhang and S. Satpathy, Phys. Rev. B **44**, 13319 (1991).]
- ¹⁸ Although we do not consider here the situation, when the insulating ground state is caused by chemical bonding effects, the extension of our arguments to this case is rather straightforward.
- ¹⁹ J. E. Hirsch, Phys. Rev. Lett. **53**, 2327 (1984).
- ²⁰ J. van den Brink, M. B. J. Meinders, J. Lorenzana, R. Eder, and G. A. Sawatzky, Phys. Rev. Lett. **75**, 4658 (1995).
- ²¹ M. B. J. Meinders, J. van den Brink, J. Lorenzana, and G. A. Sawatzky, Phys. Rev. B **52**, 2484 (1995).
- ²² J. D. Talman and W. F. Shadwick, Phys. Rev. A **14**, 36 (1976).
- ²³ J. Zaanen and O. Gunnarsson, Phys. Rev. B **40**, 7391 (1989); D. Poilblanc and T. M. Rice, *ibid.* **39**, 9749 (1989); J. Zaanen and P. B. Littlewood, *ibid.* **50**, 7222 (1994); V. I. Anisimov, I. S. Elfimov, N. Hamada, and K. Terakura, *ibid.* **54**, 4387 (1996); V. I. Anisimov, I. S. Elfimov, M. A. Korotin, and K. Terakura, *ibid.* **55**, 15494 (1997); T. Mizokawa and A. Fujimori, *ibid.* **56**, R493 (1997); T. Mizokawa and A. Fujimori, Phys. Rev. Lett. **80**, 1320 (1998).
- ²⁴ See, e.g., *ab initio* HF calculations for the semiconductors: A. Svane, Phys. Rev. B **35**, 5496

- (1987) and the transition-metal monoxides: M. D. Towler, N. L. Allan, N. M. Harrison, V. R. Saunders, W. C. Mackrodt, and E. Aprà, Phys. Rev. B **50**, 5041 (1994), in comparison with the exact KS treatment for the HF total energy functional performed for the same materials: T. Kotani and H. Akai, Phys. Rev. B **54**, 16502 (1996) and Ref. 25. Similar effect is clearly seen in *GW* calculations for semiconductors (see, e.g., Fig.1 in Ref.9).
- ²⁵ M. Städele, J. A. Majewski, P. Vogl, and A. Görling, Phys. Rev. Lett. **79**, 2089 (1997).
- ²⁶ T. Grabo, T. Kreibich, S. Kurth, and E. K. U. Gross, in *Strong Coulomb Correlations in Electronic Structure: Beyond the Local Density Approximation*, edited by V. I. Anisimov (Gordon and Breach, Tokyo, 1999).
- ²⁷ I. V. Solovyev, P. H. Dederichs, and V. I. Anisimov, Phys. Rev. B **50**, 16861 (1994).
- ²⁸ The scaling transformation naturally explains that $\Delta_{\text{HF}} \geq \Delta_{\text{KS}}$ when $v \geq 0$. In fact, a similar tendency for the KS gap and the true conductivity gap Δ was obtained by Schönhammer and Gunnarsson.¹¹ They considered a formally exact KS scheme for a Hubbard-type model, which was solved numerically for finite systems with subsequent extrapolation of results to $N \rightarrow \infty$. They found that $\Delta \approx \Delta_{\text{KS}}$ only when $v=0$, whereas more generally it held $\Delta \geq \Delta_{\text{KS}}$. They interpreted the difference between Δ and Δ_{KS} using the discontinuity scenario (see also Ref.12). Our standpoint is different.
- ²⁹ Y. Nagaoka, Phys. Rev. **147**, 392 (1966).
- ³⁰ S. H. Liu, Phys. Rev. B **15**, 4281 (1977).
- ³¹ A. I. Liechtenstein, M. I. Katsnelson, V. P. Antropov and, V. A. Gubanov, J. Magn. Magn. Matter. **67**, 65 (1987).
- ³² I. V. Solovyev and K. Terakura, Phys. Rev. B **58**, 15496 (1998).
- ³³ R. Eder, J. van der Brink, and G. A. Sawatzky, Phys. Rev. B **54**, R732 (1996).
- ³⁴ I. V. Solovyev, A. I. Liechtenstein, and K. Terakura, Phys. Rev. Lett. **80**, 5758 (1998).

TABLE I: Symmetry constraints superimposed on the density matrix and the KS potential. The uniform potential shift is set as zero. Note also that the number of electrons is $n_e = n_0 \text{Tr}(\hat{\rho}^\uparrow)$. In the COF state, the minority-spin bands are requested to be unoccupied; in all other respects the parameters $B_{1,2}^\downarrow$ can be arbitrary.

	state density matrix	KS potential
AF	$\hat{\rho}^\downarrow = \hat{\sigma}_x \hat{\rho}^\uparrow \hat{\sigma}_x$	$-B_1^\uparrow = B_1^\downarrow = B_2^\uparrow = -B_2^\downarrow$
CON	$\hat{\rho}^\downarrow = \hat{\rho}^\uparrow$	$-B_1^\uparrow = -B_1^\downarrow = B_2^\uparrow = B_2^\downarrow$
COF	$\hat{\rho}^\downarrow = 0$	$-B_1^\uparrow = B_2^\uparrow$




FIG. 1: Doubling of the unit cell associated with the spin or/and charge ordering. Two nonequivalent sites ($\nu=1,2$) are shown by white and black circles. Within a doubled unit cell, the site 1 is located in the origin and the position of the site 2 is defined by the vector $\mathbf{r}$. z is the coordination number. a is the lattice parameter. $f(\mathbf{k})$ is the structure factor - see text ($\mathbf{k}$ is in the units of $1/a$).




FIG. 2: τ_x versus τ_z for various lattices and $n_e=n_0$.




FIG. 3: Characteristic behavior of τ_x , τ_z and b calculated self-consistently as the function of the electron number near $n_e=n_0$.